

Partial characterization of *nif* genes from the bacterium *Azospirillum amazonense*

D.P. Potrich¹,
L.M.P. Passaglia^{1,3}
and I.S. Schrank^{1,2}

¹Centro de Biotecnologia, and Departamentos de
²Biologia Molecular e Biotecnologia and ³Genética,
Universidade Federal do Rio Grande do Sul, Porto Alegre, RS, Brasil

Abstract

Azospirillum amazonense revealed genomic organization patterns of the nitrogen fixation genes similar to those of the distantly related species *A. brasilense*. Our work suggests that *A. brasilense nifHDK*, *nifENX*, *fixABC* operons and *nifA* and *glnB* genes may be structurally homologous to the counterpart genes of *A. amazonense*. This is the first analysis revealing homology between *A. brasilense nif* genes and the *A. amazonense* genome. Sequence analysis of PCR amplification products revealed similarities between the amino acid sequences of the highly conserved *nifD* and *glnB* genes of *A. amazonense* and related genes of *A. brasilense* and other bacteria. However, the *A. amazonense* non-coding regions (the upstream activator sequence region and the region between the *nifH* and *nifD* genes) differed from related regions of *A. brasilense* even in nitrogenase structural genes which are highly conserved among diazotrophic bacteria. The feasibility of the 16S ribosomal RNA gene-based PCR system for specific detection of *A. amazonense* was shown. Our results indicate that the PCR primers for 16S rDNA defined in this article are highly specific to *A. amazonense* and can distinguish this species from *A. brasilense*.

Key words

- *Azospirillum amazonense*
- *nif* genes
- *glnB*
- 16S rDNA

Correspondence

I.S. Schrank
Centro de Biotecnologia
Departamento de Biologia Molecular
e Biotecnologia
Av. Bento Gonçalves, 9500
Caixa Postal 15005
91591-970 Porto Alegre, RS
Brasil

Research supported by FAPERGS
and CNPq. D.P. Potrich is the
recipient of a CAPES fellowship.

Received October 26, 2000
Accepted June 18, 2001

Introduction

Bacteria belonging to the genus *Azospirillum* are found associated with forage grasses, cereals and agriculturally important crops from various geographical origins. This genus consists of seven species, *A. lipoferum*, *A. brasilense*, *A. amazonense*, *A. halopraeferens*, *A. irakense* (1), *A. largomobile* (2), and *A. doebereineriae* (Eckert B, Baller-Weber O, Kirchof G, Halbritter A, Stoffels M and Hartmann A, unpublished results, and GenBank accession number AJ238576). Worldwide inoculation experiments using a variety of crops have shown that *Azospiril-*

lum species can contribute to the nitrogen economy (3,4). In addition, experiments with sugarcane and other crops suggest either a synergistic or an additive effect when *A. amazonense* is used in combination with other diazotrophics as inoculant (5).

A. amazonense differs in several important characteristics when compared to other *Azospirillum* species. Its nitrogenase activity has lower oxygen tolerance, its ability to use sucrose as carbon source is different, and the most remarkable difference is its sensitivity to alkaline pH (6). The genetic relationship existing between *A. amazonense* and other *Azospirillum* species was analyzed by 16S

rDNA restriction fragment length polymorphism (7). The phylogenetic analysis, based on 16S rDNA sequences, confirmed that *A. amazonense* and *A. irakense* form one cluster and that the closely related species *A. brasilense* and *A. lipoferum* form a second cluster together with *A. halopraeferens* (8).

During the last few years, nitrogen fixation (*nif*) genes have been isolated from a variety of diazotrophic organisms. In *A. brasilense* a DNA fragment of 25,100 bp encompassing the *nif* region was shown to contain the *nifHDKORFIY* (9), the *ORF2nifUSVORF4* (10), an *mcp*-like gene, the *nifENXORF3,5*, *fdxAnifQ*, and the *fixABCX* (Gross J, Vedoy C and Schrank IS, unpublished results) transcriptional units. However, to date, few genes were isolated from *A. amazonense* and none of them could be related to the biological nitrogen fixation process.

We report here the application of PCR and Southern hybridization to characterize some of the *nif* genes from *A. amazonense*. We have also analyzed the amplification products from two conserved regions within the *nifD* and *glnB* genes. Moreover, we have developed primers that specifically amplify *A. amazonense* DNA and allow us to distinguish between this species and *A. brasilense*.

Material and Methods

Bacterial strains and growth conditions

A. brasilense Sp7 (ATCC29145) and *A. amazonense* (ATCC35119) were grown on NFB and LGI minimal media, respectively, as previously described (6,11).

DNA manipulation and sequence analysis

All DNA manipulations were performed using standard techniques (12) and instructions provided by suppliers of material, enzymes or reagents. Total DNA extraction of *A. amazonense* was performed as previously described (13) and Southern blot analysis was performed with the ECL Direct Nucleic Acid Labeling and Detection Systems (Amersham Pharmacia Biotech, Uppsala, Sweden).

The nucleotide sequence determination was performed by the chain-termination method of Sanger et al. (14) using ³³P-dNTPs and the ThermoSequenase radiolabeled terminator cycle sequencing kit (Amersham). The PCR products were purified using the GFX PCR kit from Amersham. Analysis of DNA sequences and comparison with nucleotide and deduced protein sequences from other organisms were performed using the GCG (Wisconsin Package Version 9.0, Genetics Computer Group, Madison, WI, USA) computer programs (licensed to CENARGEN-EMBRAPA, Brasília, DF, Brazil).

Amplification conditions

PCR amplification of the target sequences was performed using a DNA thermal cycler (MJ Research, Waltham, MA, USA). The PCR mixture contained the reaction buffer (50 mM KCl, 10 mM Tris-HCl, pH 8.3, 2.5 mM MgCl₂), 200 μM of each dNTP, 30 pmol of each primer, 1 U of *Taq* polymerase (CenBiot enzymes, Centro de Biotecnologia, UFRGS, Porto Alegre, RS, Brazil), template DNA, and distilled water to a final concentration of 25 μl. The reaction mixture was subjected to PCR under the following conditions: heat denaturation at 94°C for 5 min and then an additional 35 cycles with heat denaturation at 94°C for 30 s, annealing (at a temperature defined for each set of primers, see Table 2) for 30 s, and DNA extension at 72°C for 30 s. After the last cycle, samples were maintained at annealing temperature for 5 min followed by 72°C for 10 min. PCR products were analyzed by gel electrophoresis (12). Primers listed in Table 2 were purchased from Oligo ETC. & Oligo Therapeutics Inc. (Wilsonville, OR, USA).

Results and Discussion

A. amazonense nif genes: hybridization and PCR

In order to understand the molecular organization of some of the *nif* genes from *A. amazonense* and to establish their relationship with the well-known *nif* genes from *A.*

brasilense, we used two different approaches. Initially, DNA fragments from different *nif* operons already characterized in *A. brasilense* were used in Southern blot hybridization to analyze the relatedness between the two *Azospirillum* species. The hybridization patterns of *A. amazonense* DNA are shown in Figure 1. The *nif* structural genes from *A. amazonense* are localized in a 6.5-kb *EcoRI* DNA fragment (Figure 1A) as previously found for *A. brasilense* genome (13). In *A. brasilense* and other diazotrophic bacteria the *nifHDK* genes are organized in a single transcriptional unit (9). To determine whether the *A. amazonense* *nif* genes are clustered in a similar manner as found in *A. brasilense* and other nitrogen-fixing organisms, we used sequences from *nif* genes (Table 1) from *A. brasilense* as DNA hybridization probes. Three different probes, which consist of the entire *nifHDK* operon (data not shown), the *nifH* gene alone (data not shown) from *A. brasilense*, and a region of the *nifD* gene (Table 1) from *A. amazonense*, showed homology with the same DNA fragment from *A. amazonense* (Figure 1A, lane 1E). This result suggests that the region comprising a 6.5-kb *EcoRI* DNA fragment probably contains the *nifHD* homologous in *A. amazon-*

ense and that the genes are organized in a single operon. Nevertheless, the hybridization pattern of the *nifD* gene amplified from *A. amazonense* revealed a different restriction pattern when *A. amazonense* and *A. brasilense* total DNA was digested with *PstI* and *SalI* restriction enzymes (Figure 1A, lanes 1P and 1S). A hybridization signal specific to *SalI* DNA fragments was obtained when *A. amazonense* total DNA was hybridized with *A. brasilense* *nifENX* and *fixABC* probes, respectively (Figure 1B, lane 1 and Figure 1C, lane 1).

In order to understand the nature of the *A. amazonense* genes that regulate nitrogen fixation we have isolated and characterized the *nifA* and *glnB* genes, which are responsible for the regulation of other *nif* genes and operons in *A. brasilense* (15,16). The presence of genes homologous to *glnB* and *nifA* was detected by hybridization of *A. amazonense* DNA with two different probes (Table 1). The entire *nifA* gene from *A. brasilense* is present within a 2.5-kb *SalI* DNA fragment (15). Heterologous hybridization revealed that the *nifA* homologue in *A. amazonense* is also found in a *SalI* DNA fragment of approximately 3 kb (Figure 1D, lane 1).

The *glnB* homologous gene was local-




Figure 1. Hybridization pattern of *Azospirillum amazonense* total DNA with *EcoRI* (1E), *PstI* (1P), and *SalI* (1S) in A, *SalI* in B, C and D, and *PstI* (1P) and *SalI* (1S) in E with the following probes: *nifD* in panel A; *nifEN* in panel B; *fixABC* in panel C; *nifA* in panel D, and *glnB* in panel E. Lane 2 represents *A. brasilense* total DNA digested with *PstI* in A, C and E, or *SalI* in B and D. Lane M contains λ HindIII as molecular marker.

ized within a 4.5-kb *Pst*I DNA fragment from *A. amazonense* (Figure 1E, lane 1P). Faint hybridization signals were detected in *A. amazonense* total DNA and also with *Pst*I-digested *A. brasilense* total DNA (Figure 1E, lane 2). Genes homologous to *glnB* have been found in *A. brasilense*, *Herbaspirillum seropedicae* and other bacteria, suggesting the presence of two copies of *glnB*-like genes in these organisms (17,18). Since no *Pst*I restriction site was found within *glnB* or *glnZ* genes from *A. brasilense* or in

the *glnB* gene from *A. amazonense*, the presence of two hybridization fragments in both strains may represent similarity between the probe and the two *glnB*-like genes.

The results presented here are the first to reveal homology between *A. brasilense* *nif* genes and *A. amazonense* total DNA. Taken together, these results suggest that the *nifHDK*, *nifENX* and *fixABC* operons and the *nifA* and *glnB* genes may be structurally homologous to *A. amazonense* counterpart genes.

The second approach to the understanding of the nature of nitrogen fixation in *A. amazonense* was based on PCR amplification of regions within the *nif/fix* genes. Several primers were selected from the sequences of the *nif/fix* genes from *A. brasilense*. The primers listed in Table 2 represent, except for *nifA*, those that revealed positive and conclusive results after PCR amplification and DNA sequencing. Three sets of primers were designed from regions within the *A. brasilense nifHDK* operon. The promoter region of this operon shows two overlapping upstream activator sequences (UAS) as the only potential NifA-binding sites (19). To date this organization seems to be unique to *A. brasilense* and may be an atypical NifA-binding site interacting with two dimers of NifA, as proposed to occur in other *nif* promoters (20). UAS primers define a region of approximately 200 bp in the *nifH* promoter sequence encompassing the two overlapping UAS. The second pair of IGR primers is derived from the intergenic region between *nifH* and *nifD* and amplifies a 310-bp region. In *A. brasilense* three inverted repeat structures have been found downstream from the *nifH* stop codon (9). Transcription analysis revealed the presence of one 1.1-kb transcript corresponding to the *nifH* gene (21). The *nifD* primer pair is specific for a 710-bp region of the *nifD* gene containing a highly conserved amino acid sequence and a serine residue which is present only in the *nifD* gene of *A. brasilense* (9).

Amplification of *A. amazonense* DNA with the above selected primers (Table 2) resulted in fragments of the predicted size (Figure 2A,B,C). All three amplified DNA

Table 1. Plasmids and probes.

Probe	Characteristics	Reference
nifHDK	6.5-kb EcoRI <i>A. brasilense</i> DNA fragment (<i>nifHDKORF1Y</i> operon)	9
nifH	2.8-kb EcoRI/PstI <i>A. brasilense</i> DNA fragment (<i>nifHDKORF1Y</i> operon)	9
nifD	<i>A. amazonense</i> PCR fragment	Present study
glnB/P _{II}	<i>A. amazonense</i> PCR fragment	Present study
nifA	2.5-kb Sall <i>A. brasilense</i> DNA fragment (<i>nifA</i> gene)	Pedrosa F (with permission)
nifEN	2.2-kb EcoRI/HindIII <i>A. brasilense</i> DNA fragment (<i>nifENX</i> operon)	Potrich D, Bressel T, Schrank IS and Passaglia L (unpublished results)
fixABC	3.0-kb PstI <i>A. brasilense</i> DNA fragment (<i>fixABCX</i> operon)	Gross J, Vedoy C and Schrank IS (unpublished results)

Table 2. Oligonucleotide primers.

Primer	Oligonucleotide sequence	Length (bp)	Amplification size (bp)	Annealing temperature
UAS1	5'CCTGACGCTGGCTCTGACGCTGG	23	200	52°C
UAS2	5'CGCGAACTGGCACGGGGATGCA	23		
IGR-up	5'ATCCCGACCCCGATCACGATG	21	310	52°C
IGR-do	5'CTTCTCGGGATACGCTTC	18		
nifD-up	5'ATCATCGGTGACTACAAC	18	710	52°C
nifD-do	5'ATCCATGTCGCGCGCAA	17		
nifA-up	5'CGCGGCGAGAGCGGCACC	18	na	42°C
nifA-do	5'GCGCCGTTGCGCAGCGG	18		
glnB-up	5'GCCATCATTAAGCCGTTCAA	20	250	56°C
glnB-do	5'AAGATCTTGCCGTCGCGGAT	20		
16S-up	5'TGATGGTTGTGGAGACTGTCAG	22	400	52°C
16S-do	5'ATTTACGCCTGACTTAAACA	21		

UAS = upstream activator sequences; IGR = intergenic region; na = no amplification; do = downstream; up = upstream.

fragments were used to probe total *A. amazonense* DNA, revealing homology with the same *EcoRI* fragment encompassing the *nifHD* genes (Figure 1A). To further characterize the PCR products, all DNA fragments were purified and sequenced. The sequences for the UAS and IGR products showed very little similarity to the *A. brasilense* counterparts (data not shown). Only the region amplified within the *nifD* gene is similar to other *nifD* gene sequences (discussed below). These results indicate that the *A. amazonense* non-coding regions described above differ from related *A. brasilense* regions even among nitrogenase structural genes which are highly conserved among diazotrophic bacteria.

A set of primers was designed (Table 2) from previously published sequences of *nifA* and *glnB* genes from *A. brasilense* (15,22). A PCR product of approximately 250 bp was detected in *A. amazonense* DNA (Figure 2D, lane 1). This product hybridized with total *A. amazonense* DNA (Figure 1E, lanes 1P and 1S) and, after DNA sequencing, revealed homology with P_{II} proteins from different organisms (discussed below). Surprisingly, no amplification product was visualized when *nifA* target primers were applied to *A. amazonense* DNA although these primers were designed on the basis of a highly conserved region found among NifA proteins from *A. brasilense*, *A. lipoferum* and related bacteria. This unexpected result suggests that the *A. amazonense nifA* gene may have greater differences in DNA sequence than found among other *nifA* genes previously characterized.

Amplification of *A. amazonense* 16S rDNA

To overcome problems with misinterpretation of the PCR results, a PCR amplification system was developed to specifically detect *A. amazonense*. The 16S rRNA sequence from *Azospirillum* species was obtained from the GenBank database. Multiple sequence alignments to other 16S rRNA gene sequences revealed one region of considerable sequence divergence from the closest relatives, as shown in Figure 3. Two primers (16S-up and 16S-do) were constructed, one based on the variable region and the other based on a lower sequence divergence region (Table 2). The PCR product generated with *A. amazonense* was 400 bp long (Figure 2E, lane 2) and was confirmed by direct sequencing (Figure 3). No PCR product was detected when *A. brasilense* DNA was the template (Figure 2E, lane 3). Using this methodology we could specifically detect and differentiate DNA templates from these two *Azospirillum* species.

Interspecies conservation of *nifgln* genes

A sequence comparison of the *A. amazonense* and *A. brasilense nifD* genes revealed a high degree of similarity at both the DNA and amino acid levels (Figure 4). One relevant difference between *A. amazonense* and *A. brasilense* NifD proteins is located at position 275 (relative to *A. brasilense* NifD sequence; 9). The *A. brasilense nifD* gene product contains a total of five cysteine residues, four of which are highly conserved




Figure 2. Specific PCR for amplification of *nif* genes from *Azospirillum amazonense*. Amplified products on 1.5% agarose gel were visualized with a UV transilluminator after ethidium bromide staining using the following primers: panel A, UAS; panel B, IGR; panel C, *nifD*; panel D, *glnB*, and panel E, rRNA from *A. amazonense* (lane 2) and *A. brasilense* (lane 3) total DNA. The negative control (lane 1) contains no template DNA in the reaction mixture. Lane M contains the 100-bp DNA ladder (Gibco/BRL).




Figure 5. *Azospirillum amazonense* glnB partial deduced amino acid sequence alignment with *A. brasilense* (GenBank accession No. X51499), *Rhodobacter sphaeroides* (GenBank accession No. AF032116), *R. capsulatus* (GenBank accession No. U25953), *Rhodospirillum rubrum* (GenBank accession No. X84158), *Azorhizobium caulinodans* (GenBank accession No. Y10213), *Rhizobium meliloti* (GenBank accession No. U50385), *Bradyrhizobium japonicum* (GenBank accession No. M26753), and *Herbaspirillum seropedicae* (GenBank accession No. U86073) P_{II} proteins. A black background indicates conserved residues in all aligned sequences, a dark grey background indicates conserved residues in at least 80% of the aligned sequences, and a light grey background indicates conserved residues in at least 6% of the aligned sequences. Multiple alignments were done with the PILEUP program, University of Wisconsin Genetics Computer Group, and the alignment editing was done with the GENDOC program considering the Dayhoff PAM 250 score table (26,27).

way of this microorganism may be similar to that of other *Azospirillum* species. The P_{II} protein from *A. amazonense* revealed a high level of similarity when compared to P_{II} from other diazotrophic bacteria (Figure 5).

The occurrence of duplicate copies of genes encoding P_{II}-like proteins now appears to be common among members of the α (*Azospirillum*) and γ (*Klebsiella*) subdivisions of the Proteobacteria class. In *A. brasilense*, P_{II} and P_Z proteins are involved differently in nitrogen-dependent regulation of various physiological functions (25). The P_{II} amino acid sequence is about 60% identical to that of P_Z and could explain the detection of two hybridization signals in *A. brasilense* and *A. amazonense* Southern blots (Figure 1E, lane 1S and lane 2).

In this report, we present distinct lines of evidence showing that the *nif* gene organization and regulation in *A. amazonense* differ,

in some aspects, from those of the best characterized *A. brasilense*. Although our results indicate the presence of counterparts of genes *nifHDK*, *nifENX*, *fixABC*, *nifA*, and *glnB* in *A. amazonense*, we found differences in restriction sites and non-coding sequences suggesting that the organization of these genes may differ from that of the related *A. brasilense*. These results support the suggestion by Fani et al. (8) that *A. amazonense* isolates may be members of a taxonomic cluster that is clearly distinct from the closely related *A. brasilense* and *A. lipoferum* species.

Acknowledgments

The authors thank J.I. Baldani for providing the bacterial strains, Dr. F. Pedrosa for providing the *nifA*-containing plasmid, and A. Schrank and S.C. da Silva for a critical reading of this manuscript.

References

- Bashan Y & Holguin G (1997). Azospirillum-plant relationships: environmental and physiological advances (1990-1996). *Canadian Journal of Microbiology*, 43: 103-121.
- Dekhil B, Cahill M, Stackebrandt E & Sly LI (1997). Transfer of *Conglomeromonas largomobilis* subsp. *largomobilis* to the genus *Azospirillum* as *Azospirillum largomobile* comb. nov., and elevation of *Conglomeromonas parooensis* subsp. *parooensis* to the new type species of *Conglomeromonas parooensis* sp. nov. *Systematic and Applied Microbiology*, 20: 72-77.
- Boddey RM & Döbereiner J (1995). Nitrogen fixation associated with grasses and cereals: Recent progress and perspectives for the future. *Fertilizer Research*, 4: 1-10.
- Bazzicalupo M & Okon Y (2000). Associative and endophytic symbiosis. In: Pedrosa F, Hungria M, Yates MG & Newton WE (Editors), *Nitrogen Fixation: From Molecules to Crop Productivity*. Kluwer Academic Publishers, London.
- Oliveira ALM, Urquiaga S, Döbereiner J & Baldani JI (2000). Biological nitrogen fixation (BNF) in micropropagated sugarcane plants inoculated with different endophytic diazotrophic bacteria. In: Pedrosa F, Hungria M, Yates MG & Newton WE (Editors), *Nitrogen Fixation: From Molecules to Crop Productivity*. Kluwer Academic Publishers, London.
- Magalhães FM, Baldani JI, Souto SM, Kuykendall JR & Döbereiner J (1983). A new acid-tolerant *Azospirillum* species. *Anais da Academia Brasileira de Ciências*, 55: 417-430.
- Grifoni A, Bazzicalupo M, Serio CD, Fancelli S & Fani R (1995). Identification of *Azospirillum* strain by restriction fragment length polymorphism of the 16S rDNA and of the histidine operon. *FEMS Microbiology Letters*, 127: 85-91.
- Fani R, Bandi C, Bazzicalupo M, Ceccherini MT, Fancelli S, Gallori E, Gerace L, Grifoni A, Miclaus N & Damiani G (1995). Phylogeny of the genus *Azospirillum* based on 16S rDNA sequence. *FEMS Microbiology Letters*, 129: 195-200.
- Passaglia LMP, Nunes CP, Zaha A & Schrank IS (1991). The nifHDK operon in the free-living nitrogen-fixing bacteria *Azospirillum brasilense* sequentially comprises genes H, D, K, a 353 bp ORF and gene Y. *Brazilian Journal of Medical and Biological Research*, 24: 649-675.
- Frazzon J & Schrank IS (1998). Sequencing and complementation analysis of the nifUSV genes from *Azospirillum brasilense*. *FEMS Microbiology Letters*, 159: 151-158.
- Döbereiner J, Marriel IE & Nery M (1976). Ecological distribution of *Spirillum lipoferum* Beijerinck. *Canadian Journal of Microbiology*, 22: 1464-1473.
- Sambrook J, Fritsch EF & Maniatis T (1989). *Molecular Cloning: A Laboratory Manual*. 2nd edn. Cold Spring Harbor Laboratory, Cold Spring Harbor, NY.
- Schrank IS, Zaha A, Araújo EF & Santos DS (1987). Construction of a gene library from *Azospirillum brasilense* and characterization of a recombinant containing the nif structural genes. *Brazilian Journal of Medical and Biological Research*, 20: 321-330.
- Sanger F, Nicklen S & Coulson AR (1977). DNA sequencing with chain-terminating inhibitors. *Proceedings of the National Academy of Sciences, USA*, 74: 5463-5467.
- Liang YY, Kaminski PA & Elmerich C (1991). Identification of a nifA-like regulatory gene of *Azospirillum brasilense* Sp7 expressed under conditions of nitrogen fixation and in the presence of air and ammonia. *Molecular Microbiology*, 5: 2735-2744.
- Arsène F, Kaminski PA & Elmerich C (1996). Modulation of NifA activity by PII in *Azospirillum brasilense*: evidence for a regulatory role of the NifA N-terminal domain. *Journal of Bacteriology*, 178: 4830-4838.
- de Zamaroczy M, Paquelin A, Peltre G, Forchhammer K & Elmerich C (1996). Co-existence of two structurally similar but functionally different PII proteins in *Azospirillum brasilense*. *Journal of Bacteriology*, 178: 4143-4149.
- Benelli EM, Souza EM, Funayama S, Rigo LU & Pedrosa FO (1997). Evidence for two possible glnB-type genes in *Herbaspirillum seropedicae*. *Journal of Bacteriology*, 179: 4623-4626.
- Passaglia LMP, Schrank A & Schrank IS (1995). The two overlapping *Azospirillum brasilense* upstream activator sequences have differential effects on nifH promoter activity. *Canadian Journal of Microbiology*, 41: 849-854.
- Morett E & Buck M (1988). NifA-dependent in vivo protection demonstrates that the upstream activator sequence of nif promoters is a protein binding site. *Proceedings of the National Academy of Sciences, USA*, 85: 9401-9405.
- de Zamaroczy M, Delorme F & Elmerich C (1989). Regulation of transcription and promoter mapping of the structural genes for nitrogenase (nifHDK) of *Azospirillum brasilense* Sp7. *Molecular and General Genetics*, 220: 88-94.
- de Zamaroczy M, Delorme F & Elmerich C (1990). Characterization of three different nitrogen-regulated promoter regions for the expression of glnB and glnA in *Azospirillum brasilense*. *Molecular and General Genetics*, 224: 421-430.
- Fu H, Hartmann A, Lowery RG, Fitzmaurice WP, Roberts GP & Burris RH (1989). Posttranslational regulatory system for nitrogenase activity in *Azospirillum* spp. *Journal of Bacteriology*, 171: 4679-4685.
- Elmerich C, Zimmer W & Vieille C (1991). Associative nitrogen fixation bacteria. In: Stacey G, Evans H & Burris R (Editors), *Biological Nitrogen Fixation*. Chapman and Hall, New York.
- de Zamaroczy M (1998). Structural homologues P₁₁ and P₂ of *Azospirillum brasilense* provide intracellular signalling for selective regulation of various nitrogen-dependent functions. *Molecular Microbiology*, 29: 449-463.
- Nicholas KB & Nicholas Jr HB (1997). GeneDoc: Analysis and visualization of genetic variation. Available at: <http://www.cris.com/ketcup/genedoc.shtml>. Accessed 1999.
- Nicholas KB, Nicholas Jr HB & Deerfield DW (1997). GeneDoc: Analysis and visualization of genetic variation. *EMB News*, 4: 14-14.